



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 06 OCT 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°.

02255556.9

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 02255556.9
Demande no:

Anmeldetag:
Date of filing: 08.08.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

UNILEVER PLC
Unilever House,
Blackfriars
London EC4P 4BQ
GRANDE BRETAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Processed tomato products and process for preparing the same

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A23L1/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

F7656 (V)

1

PROCESSED TOMATO PRODUCTS AND PROCESS FOR PREPARING THE SAME

Field of the invention

5

The present invention relates to processed tomato products and a process for preparing tomato products.

Background of the invention

10 In the industry of processing tomatoes to end products like sauces, ketchups, soups, toppings, etcetera usually two stages are distinguished: primary processing and secondary processing. Primary processing usually involves at least either hot- or cold breaking of the tomatoes and a concentration step. In the concentration step water is removed from the tomato pulp, so as to obtain a thick paste. The water removal can
15 be done by many ways, although evaporative removal of water (by heating) is the common method. The so-obtained thickened paste or puree can be stored or directly further processed into a range of finished products such as tomato sauce for pasta, tomato ketchup, etcetera.

20 Such end products generally need a specific thickness to be valued as quality products (next to a good colour, flavour, etcetera) . In order to achieve this, it is preferred that the product has (at a given percentage of soluble solids) a high consistency. Consistency in the tomato industry is often measured and expressed as Bostwick value. In the handbook "Tomato Production, Processing & Technology" (3rd
25 ed.) by W.A. Gould, CTI Publications, Timonium, Maryland, USA it is set out on page 329, 330 how Bostwick measurements on tomato purees and pastes is usually performed in the tomato processing industry and tomato research.

In part, the thickness is determined by the amount of insoluble solids present per unit
30 of tomato product. The insoluble solids are in part cellulose, pectins and other compounds that make up the structural matrix of the fruit. The amount of insolubles may vary per variety, season, growth stage, etcetera. The degree of concentration of tomato products is usually expressed in degrees Brix and is an indication of the amount of soluble solids in a tomato(product). To exemplify this: a tomato paste of 20
35 Brx is considered to be twice as much concentrated as a paste of 10 Brx of the same tomatoes.

F7656 (V)

2

Of course, in order to achieve thick products one can highly concentrate the tomato puree. The result will be a product with a high degree of Brix, a firm consistency (expressed by low Bostwick value). However, this is costly, as many kilograms of

5 tomatoes are needed to produce one kilogram of tomato product, and the evaporative concentration is also a cost factor. Furthermore, flavour and colour can be adversely affected by rigorous concentration, e.g. due to burning in the evaporators.

10 Many techniques have been developed to thicken paste without changing Brix value. Such methods include treating of the pectic substances with enzymes, adding thickeners, etcetera. These methods all have their disadvantages.

Hence, there is a need for tomato paste having a firm consistency at a reasonable
15 Brix value. Also, there is a need for processed tomato products other than tomato paste (in the strict sense of the word) which have an increased consistency.

Summary of the invention

20 It has now been found that the above may be achieved (at least in part) by a tomato paste having an increased consistency such that when measured at an insoluble solids interval of 2.5-3.6% at 12°Brix:

(1) $(\text{Bostwick value}) < 10.5 - 2.3822 \times (\text{percentage of insoluble solids})$,
when Bostwick is measured as defined in the reference given above.

25

Preferably, this is achieved by a tomato paste having an increased consistency such that when measured at an insoluble solids interval of 2.5-3.6% at 12°Brix:

(2) $(\text{Bostwick value}) < 10.0 - 2.3822 \times (\text{percentage of insoluble solids})$.

30 Most preferably, this is achieved by a tomato paste having an increased consistency such that when measured at an insoluble solids interval of 2.5-3.6% at 12°Brix:

(3) $(\text{Bostwick value}) < 9.5 - 2.3822 \times (\text{percentage of insoluble solids})$.

As is stated above, the tomato paste when measured should have an insoluble solids
35 level of 2.5-3.6%, and at 12°Brix. Pastes with different levels of Brix are also part of

F7656 (V)

3

the invention, but need to be concentrated/diluted before measurement. In the above, the Bostwick value will suitably be above 0.1 at said Brix level.

5 Tomato paste is herein to be understood as a commercially-processed (or factory-processed) tomato paste as it is known in the art of tomato processing. Such tomato paste is the result of primary processing tomatoes (comminution / heating and concentrating by removal of water) as is done shortly after harvesting. A hot break process is preferred for optimal consistency. The resulting product is a concentrated paste, which can be stored until further use, or can be sold. There are commercial
10 producers of such tomato paste (product). For comparison and measuring, such paste should be free of added thickeners, such as starches or gums. Also, for comparison and measuring, the pastes should not have been subjected to additional processing steps that may increase the consistency, such as homogenisation treatment. Conventional, commercially available paste is free of such additional
15 thickeners or process steps.

Although for measuring the Bostwick value at 12°Brix as set out above the tomato paste is a paste obtained with a hot break process, without additional process steps or ingredients that influence the consistency, the invention may be applied to all sorts
20 of tomato paste (hot and cold), which do comprise additional thickeners or process steps that influence the consistency.

Of a commercially available paste one can measure Bostwick, Brix, and insoluble solids, and such numbers can give an indication of the quality of the paste.
25

Tomato paste can be obtained by a hot break process (comminuting and heating to approx. 80°C), optionally followed by a concentration step to bring it to the required Brix value. Such concentration (i.e. water-removal) will usually be done by evaporation. The tomato paste according to the invention does not contain gums,
30 starches, or other thickeners when measuring Bostwick and Brix value. Bostwick is usually measured at 12°Brix. If the tomato paste has a too high Brix value, dilution with water to the required value of 12° may be applied.

The tomato paste according to the invention preferably is red, as expressed by
35 having a USDA color score at 8.5°Brix of 35-60. Dilution may be needed to measure at the required Brix value.

F7658 (V)

4

- There are factories ("secondary processing") that buy/use tomato paste for preparing processed tomato products, such as pasta sauce, juice, ketchup, etcetera. Such processed tomato products may also be prepared from tomato paste, or from fresh tomatoes. Following this, apart from a tomato paste with an increased consistency, there is also a desire for processed tomato products with a good consistency.

- One factor which limits the consistency that can be obtained from processed tomato products is the softening of fruit that takes place as part of the ripening process.
- 10 Aspects of fruit ripening such as development of colour and flavour give rise to desirable characteristics in processed tomato products. It would be advantageous to combine the high consistency of unripe fruit with the colour and/or flavour of ripe fruit in a processed tomato product.
- 15 A number of genes are involved in controlling the process of tomato fruit ripening. Mutations in such genes can lead to ripening-inhibited fruit in which all aspects of the ripening process, such as softening, red colour formation, and flavour development are inhibited. If the mutation is present in homozygous form, softening is minimised, and development of both colour and flavour is severely restricted. Examples of gene mutations that result in ripening-inhibition in tomatoes include 'alcobaca' (*alc*),
- 20 'ripening-inhibited' (*rin*), 'non-ripening' (*nor*), and 'Never ripe' (*Nr*).

- It has been found that both tomato paste and processed tomato products with a good consistency can suitably be achieved if tomatoes homozygous in *alc*, *rin*, *nor*, or *Nr* are used to prepare a tomato paste or processed tomato product. Hence, the invention further pertains to a paste or product comprising tomatoes which are
- 25 homozygous for *alc*, homozygous for *rin*, homozygous for *nor*, homozygous for *Nr*, heterozygous for combinations of (at least) two of the *alc*, *rin*, *nor* or *Nr* genes, or combinations thereof. Such tomatoes are herein after referred to as "tomatoes according to the invention".
- 30

- In a preferred aspect of the invention, the paste or product is prepared by using tomatoes according to the invention and which in addition comprise color enhancing genes such as old gold crimson (*ogc*), high pigment (*hp*), dark green (*dg*), intense pigment (*ip*), or color enhancing transgenic genes. Such tomatoes can not only be
- 35 used to make tomato paste but can be used for a whole range of processed tomato

F7656 (V)

5

products. The term "processed tomato product" is herein to be understood as to comprise any product that comprises tomatoes which are subjected to processing steps (in any order) such as heating and breaking and optionally concentrating and packing. Examples of processed tomato products are: tomato pastes, tomato sauces, tomato juices, tomato concentrates, tomato passatas, salsa, barbecue sauce, pizza sauce, spaghetti sauce, tomato fritto, ketchup (catsup), soup or other form.

As a result of the invention, it is possible to take advantage of the outstanding paste and serum viscosity of tomatoes which are homozygous for the *alc* genes without sacrificing desirable tomato color characteristics which are of importance to consumers. Also, the paste and serum of the tomatoes enjoy excellent resistance to syneresis. It is likewise believed that homozygous *rin* tomatoes, homozygous *nor* tomatoes, homozygous *Nr* tomatoes, or heterozygous *alc/rin*, *alc/nor*, *alc/Nr*, *rin/nor*, *Nr/nor*, *rin/Nr* tomatoes can be advantageously used in the present invention.

The tomato paste according to the invention preferably has at 12° Brix Bostwick thickness values in the range of from 0-3 cm, preferably from 0-2 cm. Likewise preferred tomato pastes according to the invention enjoy at 12° Brix syneresis levels of less than 4 mm, preferably less than 3 mm. This is in contrast to Bostwick values of 4.5-7 cm and syneresis values of 13-25 mm for, e.g., the BOS 3155 variety (industry-known variety).

The invention may provide processed tomato products having both good color and outstanding thickness, without requiring the mixing of different types of tomatoes. Preferably USDA paste color scores at 8.5° Brix for pastes of the invention are at least 35, especially greater than 42. Preferably, said color scores are below 60. USDA scores are standardised measurements for color quality.

We have found that it is possible to produce a tomato having both homozygous *alc* and the old gold crimson (*ogc*) genes, wherein the tomato color is good, yet at the same time tomato fruit firmness and juice and paste viscosity are excellent as a result of the ripening inhibiting effect of the *alc* gene.

Following the above, the present invention relates to processed tomato products such as tomato pastes, tomato sauces, tomato juices, tomato concentrates, tomato passatas, salsa, barbecue sauce, pizza sauce, spaghetti sauce, tomato fritto,

F7656 (V)

6

ketchup (catsup), soup and others, which processed tomato products comprise tomatoes according to the invention. Preferably, the above products are prepared of tomatoes which further include color enhancing genes as well. Processed tomato products preferably have a Brix value of 5-31°, preferably (depending upon the

5 intended use) of 10-25°. Also depending upon the use they may contain 0.1-5% wt, preferably 0.5-3% wt of salt, most preferably 1-2% wt. The pH may suitably be between 3 and 5, preferably between 4.0 and 4.4.

10 Preferably, the invention concerns processed tomato products made from populations or assemblages of the above fruits having an average of at least 10% by weight, and preferably at least 25%, more preferably at least 50% of the tomatoes with the above-described genes. The tomatoes for such processed tomato products may be obtained through classical breeding and selecting, but may also be obtained by genetic modification, as is set out in WO 01/04315 and WO 01/14561.

15 The pastes of the invention preferably include at least 50 % by weight of the tomatoes according to the invention, especially from 50 to 100% by weight. Juices preferably include at least 10% by weight of the tomatoes according to the invention, especially from 20 to 40% by weight.

20 Preferably, the tomatoes according to the invention are homozygous for the color enhancing gene such as *ogc*.

25 Use of the tomatoes according to the invention is particularly beneficial in view of their unique qualities, such as extremely high viscosity and almost no syneresis. It is believed these advantages are not achieved with tomatoes or tomato pastes outside of our invention (when measured with equivalent soluble solids level and absence of other thickening material, such as starch, gums, etcetera). A secondary benefit is that as a result of such characteristics, less paste can be used in preparing a sauce. The
30 advantageous paste characteristics according to the invention can be expected to translate to improved, consumer perceivable characteristics for processed tomato products, such as improved mouthfeel and texture and to lead to more full-bodied sauces and other products.

35
Detailed description of the invention

F7656 (V)

7

Although it is believed tomatoes as such are known which are homozygous in *rin* or one or more other ripening-inhibiting genes mentioned, it is believed such tomatoes have never been used in tomato processing. Also, the tomatoes homozygous in *rin* that have been studied usually referred to tomatoes that do not form color. Hence,
5 (industrially) processed tomato products (and tomato paste) having the properties as described above are novel, and in particular such processed tomato products that have a red or reddish color (e.g. USDA color score of at least 35, optionally less than 60). Furthermore, it is quite surprising that processed tomato products of good quality in terms of consistency and color could be achieved having the properties as now
10 claimed, as tomatoes that are firm are usually associated with green, unripe tomatoes. Unripe, green tomatoes are unsuitable to use in large quantities in conventional tomato products, following the color and the flavour profile which is different from ripe tomatoes.

15 Without wishing to be bound by theory, it is believed that tomatoes according to the invention are different from conventional tomatoes in that such genes may occur in conventional tomatoes, but not in a homozygous genotype. When such genes are present in homozygous form, they may interrupt part of the ripening process. It is believed (but this is speculation) that the tomatoes according to the invention have
20 different cell walls, e.g. more dense cell walls.

It has been found that a tomato believed to be homozygous in *alc* has levels of certain enzymes which are different from conventional tomatoes. It was found that such a tomato, also comprising a color gene such as *ogc*, had similar levels of
25 exogalactanase when green as a conventional tomato. This is not surprising, but several days post breaker (i.e. when pinkish/orange/red) the level of exogalactanase stayed low for the *alc/ogc* tomato, whereas for conventional tomato this level increases substantially. Regarding polygalacturonase similar findings were obtained. Processing such tomatoes into processed tomato products has distinct advantages.
30 Still, thanks to the color gene *ogc*, such tomato has good color. Although a process may involve processing only such tomatoes, it may be preferred to use a blend of tomatoes: conventional tomatoes (for economic reasons) with tomatoes according to the invention. Preferably, such tomatoes according to the invention should also have one or more of the color genes as set out hereinbefore.

35

F7656 (V)

8

Hence, the present invention also relates to a process for preparing a tomato product, the product having a USDA color score of 35-60 and wherein at least 10% (pref. at least 20%, more pref. at least 50%, up to 100%) of the tomatoes to be processed have a level of polygalacturonase of less than 200 (preferably less than

- 5 100, more pref. less than 50, usually more than 1) μ moles GalA/ml/hour, and wherein said tomatoes to be processed have a level of exogalactanase of less than 70 (preferably less than 50, most preferably less than 35, usually more than 0.1) nmoles galactose/g fwt/hour (fwt = fresh weight). In this, it is preferred that at least 10% (pref. 20%, more pref. 50%) of the tomatoes to be processed are homozygous for *rin*,
10 homozygous for *nor*, homozygous for *Nr*, homozygous for *alc*, heterozygous for combinations of two of the *rin*, *nor*, *Nr* or *alc* genes, or combinations thereof. It may also be preferred that the tomatoes are homozygous for at least two of *rin*, *nor*, *Nr*, or *alc*.
- 15 As (depending on amount used, and desired end product) it may be preferred that the resulting product has some color, it is preferred that the tomatoes as used in the process as set out above further comprise at least one color enhancing gene. For example, said color enhancing genes may be selected from the group consisting of old gold crimson (*ogc*), high pigment (*hp*), dark green (*dg*), Intense pigment (*ip*), as
20 well as color enhancing transgenic genes.

Conventional tomatoes are usually processed into paste using either a cold-break process or a hot-break process. The hot-break process involves heating to above about 80°C and comminuting ('breaking') the tomatoes, whereas a cold break would
25 be heating to below about 80°C and comminuting ('breaking') the tomatoes. The hot break process has as advantage that endogenous enzymes are inactivated quickly, including pectin-degrading enzymes like exogalactanase and polygalacturonase. Such a product (e.g. paste) with substantial amount of long pectin-chains may have good consistency. The disadvantages are that heating may involve damage to the
30 flavour: a cooked or burnt aroma may develop, losing volatiles and/or fruity flavours. The cold-break process does not de-activate the pectin-degrading enzymes quickly, so some degradation of pectin may occur, resulting in a paste with less consistency. On the other hand, the flavour is usually better of a cold break product. For these reasons, mixtures of cold- and hot break products may be used.

35

F7656 (V)

9

Following the low level of pectin-degrading enzymes (polygalacturonase and exogalactanase) of the tomatoes according to the invention, such tomatoes may be processed using a so-called 'cold break process' and have consistency more similar to hot break products, as it is believed by nature less pectin-degrading enzymes are present in the tomatoes according to the invention, and hence even if they are processed into paste using a cold-break process such tomato paste may contain a substantial amount of pectin. Thus, the invention also relates to a process comprising the steps of:

- heating tomatoes to a temperature of 60-120°C (preferably 60-80°C),

- size reduction (e.g. comminution) of said tomatoes

in any given order (optionally followed by concentration).

The tomato pastes and processed tomato products may be prepared using conventional processing techniques.

As the tomatoes according to the invention are low in certain enzymes, the invention further relates to a process for preparing a tomato product (e.g. paste or any other product) in which tomatoes are used which are low in polygalacturonase and/or exogalactanase.

Hence, the invention further relates to a process for preparing a tomato product, the product having a USDA color score of 35-60 (i.e. red) and wherein at least 10% (pref. at least 20%, more pref. at least 50%, up to 100%) of the tomatoes to be processed have a level of polygalacturonase of less than 200 (preferably less than 100, more pref. less than 50, usually more than 1) μ moles GalA/ml/hour, and wherein said tomatoes to be processed have a level of exogalactanase of less than 70 (preferably less than 50, most preferably less than 35, usually more than 0.1) nmoles galactose/g fwt/hour.

In the above, it is preferred that at least 10% (pref. at least 20%, more pref. at least 50%, up to 100%) of the tomatoes to be processed are homozygous for *rin*, homozygous for *nor*, homozygous for *Nr*, homozygous for *alc*, heterozygous for combinations of two of the *rin*, *nor*, *Nr* or *alc* genes, or combinations thereof.

The invention also relates to a process and product (i.e. tomato paste and processed tomato product) in which other ripening-inhibiting genes than *alc*, *rin*, *nor*, *Nr*, are

F7656 (V)

10

present in the tomato in such a genotypic form that they inhibit ripening similar to the tomatoes as are herein disclosed. This may relate to genes not yet known to inhibit ripening, alone or in combination.

- 5 It is preferred that the tomatoes as used in the process as set out above (e.g. to prepare a processed tomato product) further comprise at least one color enhancing gene. For example, such color enhancing genes are selected from the group consisting of old gold crimson (*ogc*), high pigment (*hp*), dark green (*dg*), intense pigment (*ip*), as well as color enhancing transgenic genes.

10

The invention further relates to a process as set out above, wherein the process comprises the steps of:

- heating tomatoes to a temperature of 60-120°C,
- comminuting or dicing said tomatoes

- 15 in any given order. More preferred in the above heating step is a temperature of 60-80°C. Optionally, a concentration step may be applied, e.g. by water-removal, e.g. by evaporation.

20 EXAMPLES

In the experiments below, the following methods were followed.

Brix

- A digital refractometer (Bellingham Stanley RFM 342 digital refractometer)
- 25 thermostatically controlled at 20°C was used. The refractometer was calibrated with a range of 1-30% w/w sucrose in de-ionised water as standard solutions. Enough tomato product was weighed into centrifuge tubes to provide a 1-2 ml liquid layer after centrifugation and centrifuged in a high speed centrifuge at 20°C, using a Beckman Optima TLX ultracentrifuge (TLA100.4 8-position fixed angle rotor) having
- 30 the following gradient: 5,000/2min., 20,000/2min., 75,000/4min., 100,000/10min., 50,000/1min., end).at 95,000 RPM +/- 5,000RPM for 5 minutes to separate liquid from solid. The centrifugate liquid was placed in a small vial and mixed gently. The liquid was placed on the optic of a thermostatted refractometer, the lid closed and measured after the sample had sat for 30 seconds to reach the required temperature.
- 35 The average of triplicate readings was calculated.

F7656 (V)

11

Bostwick

Bostwick measurement was performed on a 25cm Bostwick levelled in two directions. Paste was diluted to 12° Brix and warmed or cooled to 20°C. Sample was placed in the Bostwick to the top of the sample chamber and the trap door opened.

5 Degree of flow was determined after 30 seconds.

Samples were each tested in duplicate.

Insoluble solids

10 Tomato paste was sourced from around the world including Unilever factories in Chile (Malloa), California (Stockton/Merced), India and Australia (Tatura) and external sources (Conesa, ARC, Copais) These were used to construct a calibration line for tomato paste at 12 Brix.

15 A 1-1.5g sample of juice was weighed out to 4 decimal places between 4 pre-weighed filters (Whatman GFA 5.5cm diameter). This was then placed on a Buchner vacuum filtration system and washed with 6 litres of de-ionised water. The filters were then dried in a vacuum oven at 70° C for 1.5 hours and then cooled in a dessicator to room temperature. The filters were then re-weighed and insoluble solids calculated as the final weight minus initial filter weight, divided by the initial juice weight minus filter weight. Determination was carried out in triplicate.

20

% Insolubles = (weight of dried sample + filter) – filter/sample weight * 100

Also, the dilution factor from paste to 5°Brix juice as above was taken into account by separate multiplying.

25

Example 1: breeding and selection

As starting material a cross between a homozygous ripening-inhibited mutant (thought to be *alc*) and a homozygous old gold crimson (*ogc*) mutant was obtained from Ohio State University, USA (accession number 96-9422-400). This population of 30 F1 heterozygous *alc/ogc* was then selfed, and single plants selected that had the phenotypical characteristics of both fruit ripening-inhibition (homozygous *alc*) and golden flower colour (homozygous *ogc*). Seed from selected plants was then back-crossed with in-house breeding lines to produce stable double homozygous plants, for evaluation of fruit and processed tomato product characteristics.

35

Example 2: Bostwick vs. insoluble solids: 22 conventional pastes (control) and according to invention

Of 22 hot and cold break tomato pastes from factories or commercially available the percentage of insoluble solids was measured, and the Bostwick value was

- 5 determined (all at 12°Brix). The Bostwick values (averages of various measurements per paste) were then plotted against percentage of insoluble solids.

The hot break pastes were either commercially available products, or were prepared in own factories using the following process: the tomatoes are crushed with a minimum of air inclusion and quickly heated to greater than 85°C typically through

- 10 contact with a steam coil. The juice is then extracted and evaporated typically by a 2 or 3 stage process to between 24° and 31°Brix. For measuring, the samples were diluted to 12°Brix.

The cold break pastes were commercially available products, and are typically

- 15 prepared by crushing the tomatoes at temperatures of less than 85°C with a minimum of air inclusion. The juice is then extracted and evaporated by typically a 2 or 3 stage process to between 24 and 31°Brix. For measuring, the samples were diluted to 12°Brix.

- 20 The percentage insoluble solids was calculated as described earlier. The Bostwick value was determined using the method as described earlier. The results of these measurements are in table 1.

Table 1: Insoluble solids and Bostwick of 22 commercially available pastes

Insoluble solids	Bostwick
1.756	7.6
2.0545	6.75
2.062	6.4
2.0845	6.6
2.162	6.3
2.217	5.8
2.353	5.8
2.368	5.5
2.413	5.5
2.568 ⁽¹⁾	5.1

F7656 (V)

13

2.6515	5
2.674	4.9
2.678	4.2
2.7415	4.75
2.95 ⁽²⁾	4.5
2.974	4
2.98 ⁽³⁾	4.3
3.07	4.1
3.139	3.25
3.38	3.2
3.38	3.25
3.6 ⁽⁴⁾	2.7

Source of some of the pastes:

(1): Chilean Malloa

(2): Unilever Van den Bergh's

5 (3): CONESA

(4): COPAIS

Two trial harvests of *a/c-ogc* tomatoes (according to the invention) were processed and Bostwick/insolubles measured in the same manner, with results:

Insoluble solids	Bostwick
3.3	0.9
3.6	0.5

10

The results are set out graphically together with the results of table 1 in figure 1.

Figure 1 also gives lines for the equations:

(1) (Bostwick value) = 10.5 - 2.3822 x (percentage of insoluble solids)

15 (2) (Bostwick value) = 10.0 - 2.3822 x (percentage of insoluble solids)

(3) (Bostwick value) = 9.5 - 2.3822 x (percentage of insoluble solids)

As can be seen, all conventional pastes tested have a Bostwick value larger than equation (1) above would give, at the insoluble solids interval of 2.5-3.6%.

F7656 (V)

14

Example 3: exogalactanase activity

Exogalactanase activity of four types of tomatoes were measured at three stages of maturity: in the green stage (i.e. well before breaker point), 5-6 days post breaker,

- 5 and 12-13 days post breaker. The four types tomatoes were: the tomato according to example 1, which is thought to be homozygous for *alc* and homozygous for *ogc*, Bos 3155 (as commercially available) U338 (internal breeding line with conventional ripening and color) and homozygous *ogc* (internal breeding line with conventional ripening).

10

The exogalactanase activity was expressed as nmoles galactose per gram fresh weight (fwt) per hour that could be converted. The galactose conversion was measured using the following protocol.

15 Preparation of extracts

- Samples of tomato pericarp were taken from the frozen storage and placed in a 50ml Falcon tube containing PVPP (1% w/v buffer). 1:1.5 (w/v) of 0.2M NaPhosphate buffer pH7.5 was then added. This was left for 60 mins at 4°C for the fruit to defrost slightly to allow a more uniform homogenisation. A Polytron SEV was used for 1-2 mins to homogenise the fruit. The extract was stirred for 20 mins, left to stand for 20 mins then 20 centrifuged at 38,700 x g for 20 mins (all performed at 4°C). The supernatant was divided into 1ml aliquots for the subsequent assays and frozen at -20°C.

Exogalactanase assay

- 25 Exogalactanase activity was measured by a linked assay consisting of two steps. (1) Galactan was prepared as described [Methods in Carbohydrate Chemistry Volume 5 (pp 132-134) and incubated with extract in the presence of buffer. (2) D-galactose released in (1) was quantified by incubating with NAD and β -D-galactose dehydrogenase as described [Kurz and Wallenfels, 1974. Methods of enzymic analysis, 30 1279-1282. ed. Verlag Chemie, Weinheim].

1. The following components were mixed and incubated overnight at 30°C (in duplicate):

- | | | |
|-------|------------|-------------------------|
| Test: | 30 μ l | 10 mg/ml Lupin Galactan |
| 35 | 15 μ l | 1M Na Acetate, pH 5.0 |
| | 30 μ l | extract |

F7656 (V)

15

Control: 30 μ l H₂O
15 μ l 1M Na Acetate, pH 5.0
30 μ l extract
Substrate Control: 30 μ l 10 mg/ml Lupin Galactan
5 15 μ l 1M Na Acetate, pH 5.0
30 μ l H₂O

The reaction was stopped by incubating in a boiling water bath for 2 mins.

- 10 2. To 64 μ l of each incubated sample (above step 1) the following were added;
64 μ l Galactose Dehydrogenase [2.5 U/ml]
905 μ l 0.1M Tris/HCl pH8.6

- O.D. readings were taken at 340nm, following which 32 μ l 12.5mg/ml NAD was added.
15 This was incubated at room temperature for 1 hour and a further reading at 340nm was recorded (it was assumed that for each mole of galactose released, one mole of NADH is formed). Test Δ OD₃₄₀ (minus control and substrate control Δ ODs) was converted to nmol gal / g fwt / hr using a galactose standard curve. The results are set out in figure 2.

20

Example 4: polygalacturonase activity

Polygalacturonase activity of four types of tomato was measured at three stages of maturity: 5-6 days post breaker, and 12-13 days post breaker and 19-20 days post breaker. The four types of tomatoes as in example 3 were analysed.

25

Polygalacturonase activity was measured using the PAHBAH method [Lever, M (1972) A new reaction for colorimetric determination of carbohydrates. Anal. Biochem. 47:273-279].

30 Materials: solutions

Assay buffer stock: 50mM sodium acetate buffer, pH 4.0 with 0.2M NaCl.

Substrate: Sigma polygalacturonic acid (PGA), 0.4% stock in water (fresh)

Standard: Sigma D-galacturonic acid, 50mg/100ml stock (fresh)

- Enzyme extract: Tomato extracts were prepared as described in Example 3 except
35 that 2.5 volumes of buffer were used (in order to reduce the level of endogenous reducing sugars).

F7656 (V)

16

Megazyme fungal polygalacturonase (ammonium sulphate suspension) was used as a positive control. Enzyme was diluted 1 in 1,000 in assay buffer and stored at -20°C. Prior to use the enzyme was diluted 1 in 25 in assay buffer stock (giving 1 in 25,000 final dilution) and 10-200ul used per assay.

- 5 PAHBAH stock solution A: Slurry 10g para-hydroxybenzoic acid hydrazide (Sigma) in 60ml H₂O. Add 10 ml conc. HCl, mix and make up to 200 ml with water (pale yellow solution, may be stored in fridge for several weeks).
- PAHBAH stock solution B: Dissolve 29.4g trisodium citrate (0.05M) in 500 ml water. Add 2.2g anhydrous (2.9g dihydrate) calcium chloride (0.01M), mix well. Add 40g
- 10 NaOH (0.5M), dissolve and make up to 2 litres with water (colourless solution, may be stored in fridge for several weeks).

Materials: equipment

- Bolling water bath or dry heating block.
- 15 Temperature controlled water bath (40°C) or dry heating block.
- UV spectrophotometer and cuvettes.
- Teflon capped tubes (5ml).

Method

- 20 1. 0.25ml aliquots of 0.4% PGA were placed in teflon capped tubes.
2. 0.25ml of enzyme extract was added (final concentrations are 0.2% PGA, in 25mM Na Ac buffer pH 4.0 with 0.1M NaCl). A buffer only negative control, boiled diluted enzyme negative control and Megazyme PG positive control were included at this stage. Tubes were centrifuged briefly in a bench top microfuge (2,000 rpm) to ensure
- 25 all the assay mixture was at the bottom of the tube. Assays were started in batches of up to 24.
3. Assays were incubated at 40°C for 1 hour (start of incubation equals T0).
4. After 1 hour incubation time, 5ml of PAHBAH C (1 part of PAHBAH A to 9 parts of PAHBAH B, mixed well and kept on ice) was added to each assay and immediately
- 30 Incubated at 100°C for 6 minutes. For T0 controls, 5ml PAHBAH C was added to 0.25ml of PGA, followed by 0.25ml of appropriate enzyme dilution.
5. Samples were cooled under running water.
6. A standard curve was made (in duplicate) using galacturonic acid (0.025, 0.05, 0.1, 0.2, 0.3, 0.4 and 0.5 ml of Gal A Stock (50mg/100ml) per ml to give 59, 117, 235,
- 35 470, 705, 940 and 1175 nmoles GalA/0.5ml standards). 5mls of PAHBAH C was added, incubated at 100°C for 6 minutes then cooled under running water.

F7656 (V)

17

Gal A Stock (50mg/100ml) per ml	WATER per ml	Nmoles Gal A/0.5ml
0.5	0	1175
0.4	0.1	940
0.3	0.2	705
0.2	0.3	470
0.1	0.4	235
0.05	0.45	117
0.025	0.475	59
0	0.5	0

7. OD readings were taken at 410nm and the change in OD₄₁₀ over one hour (T₆₀ - T₀) determined. These values were multiplied by an OD to nmoles galA conversion factor (from standard curve of OD₄₁₀ vs. nmoles Gal A) and then by 10 (0.1g tissue / assay extract) to give a final value unit of nmoles galA produced / hour / g fresh weight.
- 5

The results are set out in figure 3.

F7656 (V)

18

CLAIMS

1. Tomato paste having an increased consistency such that when measured at an insoluble solids interval of 2.5-3.6% at 12°Brix:
5 (Bostwick value) < 10.5 - 2.3822 x (percentage of insoluble solids).
2. Tomato paste according to claim 1, such that when measured at insoluble solids interval of 2.5-3.6% at 12°Brix:
10 (Bostwick value) < 10.0 - 2.3822 x (percentage of insoluble solids).
3. Tomato paste according to claim 2, such that when measured at an insoluble solids interval of 2.5-3.6% at 12°Brix:
15 (Bostwick value) < 9.5 - 2.3822 x (percentage of insoluble solids).
4. Tomato paste according to claim 1-3, wherein the paste is obtained by a hot break process, and optionally followed by concentration.
5. Tomato paste according to claim 1-4, having a USDA color score at 8.5°Brix of 35-60.
- 20 6. Processed tomato product, wherein at least 10% (pref. 20%, more pref. 50%) tomatoes which are homozygous for *rin*, homozygous for *nor*, homozygous for *Nr*, homozygous for *alc*, heterozygous for combinations of two of the *rin*, *nor*, *Nr* or *alc* genes, or combinations thereof.
- 25 7. Product according to claim 6 comprising tomatoes which are homozygous for at least two genes of *rin*, *nor*, *Nr*, or *alc*.
8. Product according to claim 6-7 having USDA color scores at 8.5 Brix of at least 35.
- 30 9. Product according to claim 8 having USDA color scores of 42 or greater.
10. Product according to claim 8-9, having USDA color scores of less than 60.
- 35

11. Product according to claim 6-10 wherein said tomatoes further comprise at least one color-enhancing gene.
- 5 12. Product according to claim 11 wherein said color enhancing genes are selected from the group consisting of old gold crimson (*ogc*), high pigment (*hp*), dark green (*dg*), intense pigment (*lp*), as well as color enhancing transgenic genes.
- 10 13. Product according to claim 6-12, wherein the processed tomato product is in the form of tomato pastes, tomato sauces, tomato juices, tomato concentrates, tomato passatas, salsa, barbecue sauce, pizza sauce, spaghetti sauce, tomato fritto, ketchup (catsup), soup or other form.
- 15 14. Process for preparing a tomato product, the product having a USDA color score of 35-60 and wherein at least 10% (pref. 20%, more pref. 50%) of the tomatoes to be processed have a level of polygalacturonase of less than 200 (preferably less than 100, more pref. less than 50) μ moles GalA/ml/hour, and wherein said tomatoes to be processed have a level of exogalactenase of less than 70 (preferably less than 35) nmoles galactose/g fwt/hour.
- 20 15. Process according to claim 14, wherein at least 10% (pref. 20%, more pref. 50%) of the tomatoes to be processed are homozygous for *rin*, homozygous for *nor*, homozygous for *Nr*, homozygous for *alc*, heterozygous for combinations of two of the *rin*, *nor*, *Nr* or *alc* genes, or combinations thereof.
- 25 16. Process according to claim 14-15, wherein other ripening-inhibiting genes than *alc*, *rin*, *nor*, *Nr*, are present in the tomato in such a genotypic form that they inhibit ripening.
- 30 17. Process according to claim 14-16, wherein said tomatoes further comprise at least one color enhancing gene.
- 35 18. Process according to claim 17, wherein said color enhancing genes are selected from the group consisting of old gold crimson (*ogc*), high pigment (*hp*), dark green (*dg*), intense pigment (*lp*), as well as color enhancing transgenic genes.

F7656 (V)

20

19. Process according to claim 14-18, wherein the process comprises the steps of:

- heating tomatoes to a temperature of 60-120°C

5

- comminuting or dicing said tomatoes
in any given order.

F7656 (V)

21

ABSTRACT

Processed tomato products which have an increased consistency as expressed by reduced Bostwick at certain insoluble solids level and Brix value.

5

Bostwick at 12 Brix vs insolubles solids




Figure 1

2/3




Figure 2: exogalactanase levels in several tomatoes.

3/3




Figure 3: Polygalacturonase activity in several tomatoes.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.